arXiv:0912.4582v2 [quant-ph] 30 Mar 2010

ON NONLINEAR COHERENT STATES OF TRAPPED IONS

Francesco A. Raffa,[1] Mario Rasetti,[2] and Marco Genovese[3]
[1] *Dipartimento di Meccanica, Politecnico di Torino, 10129 Torino, Italy*
[2] *Dipartimento di Fisica & Scuola di Dottorato,*
Politecnico di Torino, 10129 Torino, Italy
ISI Foundation, 10133 Torino, Italy
[3] *INRIM, I-10135, Torino, Italy*

(Dated: March 31, 2010)

In order to provide experimentalists with a useful tool for experiments on the k-boson nonlinear coherent states, we investigate some critical features of these states, as regards their realization as stationary states of a trapped laser-driven ion. After illustrating the k-boson case, we analyze the single-boson nonlinear coherent states and prove that their coherence and minimum-uncertainty properties are violated whenever the Lamb-Dicke parameter is one of the roots of the Laguerre polynomials which enter such states. We discuss the possibility of circumventing this problem through the relaxing of the rotating wave approximation.

PACS numbers: 03.65.-w, 42.50.Dv, 03.65.Fd

I. INTRODUCTION

Coherent states have become a tool of the utmost importance in quantum optics and many efforts have been devoted to investigate them, also with respect to their prospective applications in the growing field of quantum technologies (QT) such as quantum information (coding, transmission and elaboration of information by exploiting properties of quantum states), quantum computing, quantum imaging and quantum metrology.

In particular, their original definition has been generalized, so that nowadays several quantum optical extensions of the Glauber states have been constructed and analyzed from both the theoretical [1]-[14] and the experimental [15]-[18] point of view. Such states can find application also in quantum calculus, as shown, e.g., in [9].

In the present paper the focus is on the k-boson and single-boson nonlinear coherent states (NLCS) which can be realized experimentally by means of trapped ions interacting with driving electromagnetic fields, as reported in references [7] and [19], respectively. Due to their mathematical structure these states are affected, for specific values of the Lamb-Dicke parameter, by singularity problems which do not appear to have been tackled systematically so far. The paper is organized as follows: in section 2 the Hamiltonian of a ion trap driven by two lasers is illustrated, while the construction of the corresponding k-boson NLCS is reported in section 3 resorting to standard tools of quantum optics, i.e., the interaction representation and the rotating wave approximation (RWA). Sections 4 and 5 are devoted to the special case of the single-boson NLCS. The relaxing of the RWA in order to remove the singularity problems is discussed in section 6.

II. THE PHYSICAL SYSTEM

The properties of the ion-trap systems and of their Hamiltonians in various physical regimes have been widely investigated [20]-[29]. As for the construction of coherent states for ion-trap systems, in [7] it has been proved that single-boson NLCS can be realized by exploiting the Hamiltonian $H = H_0 + H_{\rm int}$ of a two-level ion trapped in an external harmonic oscillator potential and interacting with two laser fields.

The free Hamiltonian H_0 describes the motion of both the electronic (or internal) and vibrational (or external) degrees of freedom of the ion, while the interaction Hamiltonian $H_{\rm int}$ corresponds to the interaction of the ion with the electromagnetic fields of the driving lasers. The external states of the ion are the ground state $|g\rangle$ and the excited state $|e\rangle$, their transition frequency is $\omega = \omega_e - \omega_g$, ν is the frequency of the trap harmonic potential. With a the annihilation operator for the vibrational motion of the trapped ion, $\hat{n} = a^\dagger a$, σ_z the Pauli matrix, which acts as a third component pseudospin operator,

$$H_0 \,=\, \hbar\nu\hat{n} + \frac{1}{2}\hbar\omega\sigma_z \,. \tag{1}$$

In Eq. (1) the zero energy level of the ion has been set at halfway between $|g\rangle$ and $|e\rangle$, so that in outer product notation $\sigma_z = |e\rangle\langle e|$ - $|g\rangle\langle g|$. The interaction Hamiltonian reads

$$H_{\rm int} \,=\, \lambda \left\{ E_0\, e^{i(k_0 x - \omega_0 t)} + E_1\, e^{i(k_1 x - \omega_1 t)} \right\} S_+ + \text{H.c.}\,. \tag{2}$$

In Eq. (2) λ is the dipole coupling matrix element, E_0, E_1, and ω_0, ω_1 are the amplitudes and the frequencies of the lasers fields, k_0, k_1 their wavectors; $x = \eta(a^\dagger + a)/k_L$ is the position operator of the ion center of mass, where $k_L \approx k_0 \approx k_1$ is the value of both lasers wavevectors, $\eta = k_L\sqrt{\hbar/(2M\nu)}$ is the Lamb-Dicke parameter, with M the ion mass. With σ_x and σ_y Pauli matrices, $S_+ = (\sigma_x + i\sigma_y)/2 = |e\rangle\langle g|$ and $S_- = S_+^\dagger$ are the two-level electronic flip operators. For the lasers tunings one sets $\omega_0 = \omega$, which corresponds to the resonant condition of the first laser, and $\omega_1 = \omega - k\nu$, which means that the second laser is tuned on k-th lower (or red-sideband) vibrational level, $k \geq 1$. From Eq. (2) one then obtains

$$H_{\rm int} \;=\; \lambda e^{-i\omega t}\left[E_0\, e^{i\eta(a^\dagger + a)} + e^{ik\nu t}E_1\, e^{i\eta(a^\dagger + a)}\right] S_+ + \text{H.c.}\;. \tag{3}$$

III. THE k-BOSON NLCS

The underlying idea for the construction of nonlinear bosonic coherent states from the Hamiltonian (3) can be summarized as follows.

1. Through the unitary transformation

$$U(t) \;=\; \exp\left(-iH_0t/\hbar\right) \;=\; \exp\left(-i\nu\hat{n}t\right)\exp\left(-i\omega\sigma_z t/2\right)\;,$$

one writes the Hamiltonian (3) in the interaction representation $\mathcal{H}_{\rm int} = U^\dagger(t)H_{\rm int}U(t)$; utilizing the RWA $\mathcal{H}_{\rm int}$ is put in the form [19]

$$\mathcal{H}^{(0)}_{\rm int} \propto \left(\hat{F}^{(0)}_k\, S_+ + \text{H.c.}\right)\;, \tag{4}$$

where $\hat{F}^{(0)}_k$ is a non-Hermitian operator acting in the Fock space;

2. the condition $\hat{F}^{(0)}_k|\xi_k\rangle = 0$ [6],[19] determines the states $|\xi_k\rangle$ upon expanding them in terms of number states; besides, $\langle\xi_k|\mathcal{H}^{(0)}_{\rm int}|\xi_k\rangle = 0$ by construction;

3. the states $|\xi_k\rangle$ are recognized to be the eigenstates of the annihilation operator

$$A_k \;=\; f(\hat{n})a^k\;, \tag{5}$$

with the appropriate identification of the nonlinear function $f(\hat{n})$, i.e., they are coherent states in the Barut-Girardello sense [30].

Let us detail the above steps. In the interaction representation one obtains

$$\begin{aligned}\mathcal{H}_{\rm int} \;&=\; \lambda E_1 e^{-\eta^2/2}\left[\frac{E_0}{E_1}\sum_{\ell,m=0}^{\infty}\frac{(i\eta)^{m+\ell}}{m!\ell!}e^{i\nu t(m-\ell)}{a^\dagger}^m a^\ell\right.\\ &\left.+\;\sum_{\ell,m=0}^{\infty}\frac{(i\eta)^{m+\ell}}{m!\ell!}e^{i\nu t(m-\ell+k)}{a^\dagger}^m a^\ell\right] S_+ + \text{H.c.}\;.\end{aligned} \tag{6}$$

The link between the physical system of the laser-driven ion trap and its mathematical structure is more clearly grasped rewriting Eq. (6) in the form

$$\mathcal{H}_{\rm int} \;=\; \lambda E_1 e^{-\eta^2/2}\left[\frac{E_0}{E_1}\,\hat{O} + \hat{Q}_k\right] S_+ + \text{H.c.}\;, \tag{7}$$

where $\hat{O}$ and $\hat{Q}_k$ are the operators

$$\hat{O} \;=\; \sum_{r=1}^{\infty} {a^\dagger}^r\,\frac{(i\eta)^r e^{i\nu rt}L^{(r)}_{\hat{n}}(\eta^2)}{(\hat{n}+1)\cdots(\hat{n}+r)} + \frac{(i\eta)^r e^{-i\nu rt}L^{(r)}_{\hat{n}}(\eta^2)}{(\hat{n}+1)\cdots(\hat{n}+r)}\,a^r + L^{(0)}_{\hat{n}}(\eta^2)\;, \tag{8}$$

$$\hat{Q}_k = \sum_{s=1}^{\infty} \frac{(i\eta)^{s+k} e^{-i\nu st} L_{\hat{n}}^{(s+k)}(\eta^2)}{(\hat{n}+1)\cdots(\hat{n}+s+k)}\, a^{s+k} + \sum_{j=1}^{k} \frac{(i\eta)^j e^{i\nu(k-j)t} L_{\hat{n}}^{(j)}(\eta^2)}{(\hat{n}+1)\cdots(\hat{n}+j)}\, a^j$$
$$+ \; e^{i\nu kt} L_{\hat{n}}^{(0)}(\eta^2) + \sum_{s=k+1}^{\infty} a^{\dagger\, s-k}\, \frac{(i\eta)^{s-k} e^{i\nu st} L_{\hat{n}}^{(s-k)}(\eta^2)}{(\hat{n}+1)\cdots(\hat{n}+s-k)} \; . \tag{9}$$

In Eqs. (8), (9), $L_{\hat{n}}^{(\alpha)}(\eta^2)$, with α integer ≥ 0, is the 'Laguerre polynomial operator'. By this we means an appropriate integral representation [31]. Its action in the Fock space results into the generalized Laguerre polynomial of degree n and parameter α [32]

$$L_n^{(\alpha)}(x) = \sum_{\ell=0}^{n} (-)^{\ell} \binom{n+\alpha}{n-\ell} \frac{x^{\ell}}{\ell!} \; . \tag{10}$$

Applying the RWA to the interaction Hamiltonian (7) implies that all terms oscillating with frequency ν or larger than ν are disregarded in Eqs. (8) and (9) so that in this approximation $\hat{O}$ and $\hat{Q}_k$ reduce to

$$\hat{O}^{(0)} = L_{\hat{n}}^{(0)} \quad , \quad \hat{Q}_k^{(0)} = (i\eta)^k \prod_{j=1}^{k} (\hat{n}+j)^{-1} L_{\hat{n}}^{(k)}(\eta^2) a^k \; .$$

This leads to the form (4)

$$\mathcal{H}_{\text{int}}^{(0)} = \lambda E_1 e^{-\eta^2/2} \hat{F}_k^{(0)} S_+ + \text{H.c.} \; , \tag{11}$$

where

$$\hat{F}_k^{(0)} = \frac{E_0}{E_1} \hat{O}^{(0)} + \hat{Q}_k^{(0)} \; . \tag{12}$$

The states $|\xi_k\rangle$, which are the stationary states of the master equation of the system, are found from the condition $\hat{F}_k^{(0)}|\xi_k\rangle = 0$. Their physical meaning is twofold: they are both noiseless codes in the sense specified in [33], as $\langle\xi_k|\mathcal{H}_{\text{int}}^{(0)}|\xi_k\rangle = 0$, and the coherent states of the system, defined as the eigenstates of the operator (5)

$$A_k \, |\xi_k\rangle = k \, \xi_k \, |\xi_k\rangle \; , \tag{13}$$

where

$$\xi_k = -\left(-\frac{i}{\eta}\right)^k \frac{E_0}{E_1} \; , \tag{14}$$

while the nonlinear operator $f(\hat{n})$ is

$$f(\hat{n}) = \prod_{j=1}^{k} (\hat{n}+j)^{-1} \, \frac{L_{\hat{n}}^{(k)}(\eta^2)}{L_{\hat{n}}^{(0)}(\eta^2)} \; . \tag{15}$$

One finds that the following self-splitting into k states results from the condition $\hat{F}_k^{(0)}|\xi_k\rangle = 0$ with $|\xi_k\rangle = \sum_{n=0}^{\infty} c_n^{(k)}|n\rangle$

$$|\xi_k\rangle = \sum_{\ell=0}^{k-1} |\xi_k\rangle_{\ell} \; . \tag{16}$$

The states $|\xi_k\rangle_\ell$,

$$|\xi_k\rangle_\ell = \sum_{n=0}^{\infty} c_{nk+\ell}^{(k)} |nk+\ell\rangle \; , \tag{17}$$

are fixed by the corresponding sets of coefficients

$$c_{nk+\ell}^{(k)} \;=\; \frac{\sqrt{\ell!}\,\xi_k^n}{\sqrt{(nk+\ell)!}} \left[\prod_{m=0}^{n-1} f(\ell+mk)\right]^{-1} c_\ell^{(k)} \;, \tag{18}$$

$c_\ell^{(k)}$ being the normalization constants, chosen to be real to retrieve the Glauber states when $k=1$ and $f \equiv 1$

$$c_\ell^{(k)} \;=\; \left\{\sum_{n=0}^{\infty} \frac{\ell!\,|\xi_k|^{2n}}{(nk+\ell)!} \left[\prod_{m=0}^{n-1} f(\ell+mk)\right]^{-2}\right\}^{-\frac{1}{2}} \;. \tag{19}$$

In Eqs. (18), (19) the scalar functions $f(\ell+mk)$ are obtained from the action of the operator (15) in the Fock space, while $\prod_{m=0}^{n-1} \doteq 1$ for $n=0$.

It is readily verified that the states (17) are eigenstates of the operator (5), $A_k|\xi_k\rangle_\ell = \xi_k|\xi_k\rangle_\ell$, with the k-degenerate eigenvalue (14); since ${}_j\langle\xi_k|\xi_k\rangle_\ell = \delta_{j\ell}$, they form an orthonormal basis of a k-dimensional Hilbert space. On the other hand, due to the splitting (16), $|\xi_k\rangle$ is eigenstate of A_k with the nondegenerate eigenvalue $k\xi_k$.

IV. THE SINGLE-BOSON NLCS

In the nondegenerate case $k=1$ one retrieves the construction reported in [7]. The notation adopted so far can be simplified dropping the index k throughout.

A. The states $|\xi\rangle$ as eigenstates of $A = f(\hat{n})a$

Eqs. (13) and (14) show that $|\xi\rangle$ is by construction eigenstate of the operator (5) with $k=1$, $A = f(\hat{n})a$, the eigenvalue being $\xi = iE_0/(\eta E_1)$. The relevant commutation relations are $[\hat{n}, A] = -A$, $\left[\hat{n}, A^\dagger\right] = A^\dagger$ and

$$\left[A, A^\dagger\right] \;=\; (\hat{n}+1)f^2(\hat{n}) - \hat{n}f^2(\hat{n}-1) \doteq C(\hat{n}) \;. \tag{20}$$

In terms of number states

$$|\xi\rangle \;=\; \mathcal{N}\sum_{n=0}^{\infty} c_n|n\rangle \;, \tag{21}$$

where the normalization constant $\mathcal{N}$, which is the new label of $c_0^{(1)}$ in Eq. (19), has been factorized so that from Eq. (18)

$$c_n \;=\; \frac{\xi^n}{\sqrt{n!}} \prod_{j=0}^{n-1} [f(j)]^{-1} \;=\; \xi^n\sqrt{n!} \prod_{j=0}^{n-1} \frac{L_j^{(0)}(\eta^2)}{L_j^{(1)}(\eta^2)} \;, \tag{22}$$

with $c_0 \doteq 1$, $f(j) = L_j^{(1)}(\eta^2)/[(j+1)\,L_j^{(0)}(\eta^2)]$ and

$$\mathcal{N} \;=\; \left[\sum_{n=0}^{\infty} |c_n|^2\right]^{-\frac{1}{2}} \;. \tag{23}$$

With $m \geq 1$, the following recurrence formula for the coefficients (22) holds

$$c_{n+m} \;=\; \sqrt{\frac{n!}{(n+m)!}} \left[\prod_{\ell=0}^{m-1} f(n+\ell)\right]^{-1} \xi^m c_n \;. \tag{24}$$

B. The states $|\xi\rangle$ as minimum-uncertainty states

For the dimensionless operators $\hat{X} = (A^\dagger + A)/\sqrt{2}$, $\hat{P} = i(A^\dagger - A)/\sqrt{2}$, in view of Eq. (20), one has $[\hat{X}, \hat{P}] = iC(\hat{n})$, while the variances

$$(\Delta\hat{X})^2 \;=\; \frac{1}{2}\left[(\Delta A)^2 + (\Delta A^\dagger)^2 + 2\langle A^\dagger A\rangle - 2|\langle A\rangle|^2 + \langle C(\hat{n})\rangle\right] \;, \tag{25}$$

$$(\Delta\hat{P})^2 \;=\; -\frac{1}{2}\left[(\Delta A)^2 + (\Delta A^\dagger)^2 - 2\langle A^\dagger A\rangle + 2|\langle A\rangle|^2 - \langle C(\hat{n})\rangle\right] \;, \tag{26}$$

calculated in the coherent states $|\xi\rangle$, minimize the Heisenberg uncertainty relation $(\Delta\hat{X})^2(\Delta\hat{P})^2 \geq |\langle[\hat{X}, \hat{P}]\rangle|^2/4$

$$(\Delta\hat{X})^2_{|\xi\rangle} \;=\; (\Delta\hat{P})^2_{|\xi\rangle} \;=\; \frac{1}{2}\,|\langle\xi|C(\hat{n})|\xi\rangle| \;. \tag{27}$$

In Eq. (27) $\langle\xi|C(\hat{n})|\xi\rangle = \mathcal{N}^2\sum_{n=0}^{\infty}|c_n|^2\,C(n) = \sum_{n=0}^{\infty}p_nC(n)$, where $C(n) = (n+1)f^2(n) - nf^2(n-1)$ and $p_n = |c_n|^2\mathcal{N}^2$ with $0 \leq p_n \leq 1\ \forall\, n$. Eq. (27) confirms that the states (21) are intelligent states along with the Glauber states. On the other hand they are not coherent states according to the Perelomov definition [34] since, in view of (20), A and $A^\dagger$ generate an infinite-dimensional algebra.

V. THE SINGULARITIES

Eq. (22) shows that for $n \geq 1$ c_n has $n(n-1)/2$ zeros and as many poles due to the roots of the n Laguerre polynomials $L_j^{(0)}(\eta^2)$ and $L_j^{(1)}(\eta^2)$, respectively, whose effects on the coherent states (21) have to be investigated.

It is worth acknowledging that the above problems are referred to briefly in [23] where it is stated that the values of η^2 must be such as to prevent them. Furthermore, in [24] the effects of an ill-defined function $f(j)$ are investigated in relation to the representation of the Fock states in terms of single-boson NLCS.

A. A single zero of c_n

Let us investigate the effect produced by a single zero, $\eta^2 = \bar{\eta}^2$, of a given c_n, say c_β. This means that $L_{\beta-1}^{(0)}(\bar{\eta}^2) = 0$ and, equivalently, $f(\beta-1) \to \infty$. Since $c_\beta = 0$, from Eq. (24) it follows that $c_{\beta+m} = 0$, $\forall\ m \geq 1$. Due to the above conditions the sums in Eqs. (21) and (23) can be truncated so that the states and their normalization constant read

$$|\xi\rangle_\beta \;=\; \mathcal{N}_\beta\sum_{n=0}^{\beta-1}c_n|n\rangle \;, \tag{28}$$

$$\mathcal{N}_\beta \;=\; \left[\sum_{n=0}^{\beta-1}|c_n|^2\right]^{-\frac{1}{2}} \;. \tag{29}$$

The vanishing of the coefficient c_β violates the coherence of the states (28) as the latter do not satisfy the eigenvalue equation of the annihilation operator

$$A\,|\xi\rangle_\beta \;=\; \xi\left(|\xi\rangle_\beta - \mathcal{N}_\beta c_{\beta-1}|\beta-1\rangle\right) \;.$$

Notice that the truncactions (28) and (29) do not remove the singularity of the states which influences the variances of the observables $\hat{X}$, $\hat{P}$. In fact, addressing now the minimum-uncertainty property in the states $|\xi\rangle_\beta$, from

$$(\Delta A)^2_\beta \;=\; \xi^2\left[\mathcal{N}_\beta^2\,|c_{\beta-1}|^2 - \mathcal{N}_\beta^4\,|c_{\beta-1}|^4 - \mathcal{N}_\beta^2\,|c_{\beta-2}|^2\right] \;,$$

and from Eqs. (25), (26), one gets $(\Delta\hat{X})^2_\beta = \Lambda + \Xi$, $(\Delta\hat{P})^2_\beta = -\Lambda + \Xi$, where

$$\Lambda \;=\; \frac{1}{2}\left(\xi^2 + \xi^{*2}\right)\left[p_{\beta-1}(\beta) - p^2_{\beta-1}(\beta) - p_{\beta-2}(\beta)\right] \;,$$

$$\Xi \;=\; |\xi|^2p_{\beta-1}(\beta)\left[1 - p_{\beta-1}(\beta)\right] + \frac{1}{2}\sum_{n=0}^{\beta-1}p_n(\beta)C(n) \;,$$



FIG. 1: The product $(\Delta\hat{X})^2_\beta\ (\Delta\hat{P})^2_\beta$ for $\beta = 4$, $|\xi| = 0.1$

while $p_n(\beta) = |c_n|^2\mathcal{N}^2_\beta$, $0 \leq p_n(\beta) \leq 1\ \forall\, n, \beta$ and $*$ denotes complex conjugation.

As an example, in Fig. 1 the product $(\Delta\hat{X})^2_\beta\ (\Delta\hat{P})^2_\beta = \Xi^2 - \Lambda^2$ is plotted in the range $0 < \eta^2 \leq 5$ assuming that the singularity problem corresponds to $\beta = 4$, i.e., $\bar{\eta}^2$ is one of the roots of $L_3^{(0)}(\eta^2)$. Of course, in Fig. 1, η^2 varies with continuity so that the product of the variances contain the roots in the considered range of all the Laguerre polynomials involved in the $\beta = 4$ case, i.e., $L_j^{(0)}(\eta^2)$ and $L_j^{(1)}(\eta^2)$, $j = 1, 2, 3$. One emphasizes that a root of the polynomials $L_j^{(0)}(\eta^2)$ (the single zero case) corresponds to a singularity of the product $(\Delta\hat{X})^2_\beta\ (\Delta\hat{P})^2_\beta$, while when a root of the polynomials $L_j^{(1)}(\eta^2)$ occurs (the single pole case of Sect. 5.2) the product vanishes.

Notice also that in the example considered the minimum-uncertainty product 1/4 is still maintained over a wide range of values of η^2. This feature strongly depends on the parameter β, in the sense that the lower the value of β, the wider the ranges where, due to the minimum-uncertainty property, the truncated states (28) can still be considered, approximately, intelligent states and conveniently utilized for QT purposes.

B. A single pole of c_n

Here $\bar{\eta}^2$ is a single pole of a given c_μ, i.e., $c_\mu \to \infty$. This implies that $L_{\mu-1}^{(1)}(\bar{\eta}^2) = 0$ and correspondingly $f(\mu - 1) = 0$. From the recurrence relation (24) it follows that $c_{\mu+m} \to \infty$, $\forall\ m \geq 1$, as rapidly as c_μ. The above conditions imply that the contribution of the first μ terms to the sums (21) and (23) can be disregarded with respect to the singular terms, leading to the following expressions of the states and their normalization constant

$$|\xi\rangle_\mu\ = \mathcal{N}_\mu \sum_{n=\mu}^{\infty} c_n |n\rangle\ , \tag{30}$$

$$\mathcal{N}_\mu\ = \left[\sum_{n=\mu}^{\infty} |c_n|^2\right]^{-\frac{1}{2}}\ . \tag{31}$$

Extracting the singular coefficient c_μ from both Eqs. (30) and (31) and utilizing the recurrence property (24), one finds

$$|\xi\rangle_\mu \;=\; [S(\mu)]^{-\frac{1}{2}} \sum_{n=\mu}^{\infty} \frac{\xi^n}{\sqrt{n!}} \prod_{\ell=\mu}^{n-1} f^{-1}(\ell)|n\rangle$$

$$\;=\; [S(\mu)]^{-\frac{1}{2}} \sum_{j=0}^{\infty} \frac{\xi^{\mu+j}}{\sqrt{(\mu+j)!}} \prod_{\ell=\mu}^{\mu+j-1} f^{-1}(\ell)|n\rangle \;,$$

where $S(\mu)$ is the series

$$S(\mu) \;=\; \sum_{j=0}^{\infty} \left[\frac{|\xi|^{2(\mu+j)}}{(\mu+j)!} \prod_{\ell=\mu}^{\mu+j-1} f^{-2}(\ell) \right]^2_{\eta^2=\bar{\eta}^2} \;, \tag{32}$$

The question whether or not the normalization constant $S(\mu)$ converges, and hence the states (30) exist (in the sense that they are normalizable), can be tackled, for example, through the ratio test. According to this criterion the convergence of the series (32), written as $S(\mu) = \sum_{j=0}^{\infty} s_j^{(\mu)}$, implies the evaluation of the ratio

$$\frac{s_{j+1}^{(\mu)}}{s_j^{(\mu)}} \;=\; \frac{|\xi|^2}{(\mu+j+1)f^2(\mu+j)} \;=\; |\xi|^2(\mu+j+1) \left[\frac{L_{\mu+j}^{(0)}(\eta^2)}{L_{\mu+j}^{(1)}(\eta^2)} \right]^2_{\eta^2=\bar{\eta}^2} \;,$$

when $j \to \infty$. Resorting to the asymptotic formulas for the Laguerre polynomials of large degree reported, e.g., in [35], one finds that

$$\lim_{j\to\infty} \frac{s_{j+1}^{(\mu)}}{s_j^{(\mu)}} = |\xi|^2 \lim_{j\to\infty} \left[\frac{\bar{\eta}}{\tan\left(2\sqrt{(\mu+j)\bar{\eta}^2} - \frac{\pi}{4}\right)} \right]^2 \;. \tag{33}$$

The result (33) proves to be wildly oscillating as j increases, which means that the very definition of the states $|\xi\rangle_\mu$ is questionable when a pole of a coefficient c_μ is encountered. This pathology becomes manifest in the example of Fig. 1 where the poles correspond to the vanishing of $(\Delta\hat{X})^2_\beta\,(\Delta\hat{P})^2_\beta$.

The above analysis shows that there are values of η^2 which alter significantly the structure of the single-boson NLCS, resulting either in the truncated states (28) or in the states (30) whose definition is not certain. These singularities are inherent in the very construction of such states which relies on the application of the RWA to the two-level ion model. One is therefore led to investigate whether relaxing the RWA can remove the above difficulties.

VI. RELAXING THE RWA

A detailed analysis of the possible corrections of the RWA is out of the aims of the present paper. Here we discuss two ways through which a 'first-order' correction of the RWA can be implemented, where first-order means that only the terms of Eqs. (8), (9) associated to the exponentials $\exp(\pm i\nu t)$ are retained. Specifically, one adds the following time-dependent operator $\mathcal{H}_{\rm int}^{(1)}$ to the interaction Hamiltonian (11)

$$\mathcal{H}_{\rm int}^{(1)} \;=\; \lambda E_1 e^{-\eta^2/2} \left[\frac{E_0}{E_1}\hat{O}^{(1)} + \hat{Q}_k^{(1)} \right] S_+ + \text{H.c.} \;,$$

where

$$\hat{O}^{(1)} \;=\; (i\eta) \left[e^{i\nu t} a^\dagger \frac{L_{\hat{n}}^{(1)}(\eta^2)}{\hat{n}+1} + \frac{L_{\hat{n}}^{(1)}(\eta^2)}{\hat{n}+1} a e^{-i\nu t} \right] \;, \tag{34}$$

$$\hat{Q}_k^{(1)} \;=\; \frac{(i\eta)^{k+1} L_{\hat{n}}^{(k+1)}(\eta^2)\, e^{-i\nu t}}{(\hat{n}+1)\cdots(\hat{n}+k+1)} a^{k+1} + \frac{(i\eta)^{k-1} L_{\hat{n}}^{(k-1)}(\eta^2)\, e^{i\nu t}}{(\hat{n}+1)\cdots(\hat{n}+k-1)} a^{k-1}$$

$$\;+\; \delta_{k,1} e^{i\nu k t} L_{\hat{n}}^{(0)}(\eta^2) \;. \tag{35}$$

The first approach develops along the general scheme of section 3, i.e., one assumes that, in this approximation, the coherent states $|z\rangle$,

$$|z\rangle \;=\; \sum_{n=0}^{\infty} d_n |n\rangle \;,$$

are determined from the condition $\hat{\mathcal{F}}|z\rangle = 0$, $\hat{\mathcal{F}} = \hat{F}_1^{(0)} + \hat{F}_1^{(1)}$ with $\hat{F}_1^{(0)}$ from Eq. (12) and $\hat{F}_1^{(1)} = (E_0/E_1)\,\hat{O}^{(1)} + \hat{Q}_1^{(1)}$, where $\hat{O}^{(1)}$ and $\hat{Q}_1^{(1)}$ are given in Eqs. (34) and (35). The coefficients d_n can be calculated from the relations

$$\frac{(i\eta)^2 e^{-i\nu t}}{\sqrt{2}} d_2 + i\eta e^{-i\nu t} w(t) d_1 + w(t) d_0 \;=\; 0 \;, \tag{36}$$

$$\begin{aligned} &\frac{(i\eta)^2 e^{-i\nu t}}{\sqrt{(n+1)(n+2)}} L_n^{(2)}(\eta^2) d_{n+2} + \frac{i\eta e^{-i\nu t} w(t)}{\sqrt{n+1}} L_n^{(1)}(\eta^2) d_{n+1} \\ &+ w(t) L_n^{(0)}(\eta^2) d_n + \frac{E_0}{E_1}\frac{i\eta e^{i\nu t}}{\sqrt{n}} L_{n-1}^{(1)}(\eta^2) d_{n-1} \;=\; 0 \;, \quad n \geq 1 \;, \end{aligned} \tag{37}$$

where $w(t) = E_0/E_1 + \exp(i\nu t)$, if d_0 and d_1 are known. To fix the latter coefficients one could resort, e.g., to suitable orthonormality conditions.

The second approach entails the utilization of the RWA according to the procedure proposed in [7]: one defines $\hat{G}_1^{(0)} = \dfrac{E_0}{E_1}\displaystyle\sum_{\ell=1}^{\infty}\frac{(i\eta)^{2\ell}}{(\ell!)^2} a^{\dagger\ell} a^{\ell} + i\eta \frac{L_{\hat{n}}^{(1)}(\eta^2)}{\hat{n}+1} a = \hat{F}_1^{(0)}$ - E_0/E_1, so that $\hat{G}_1^{(0)}|\xi\rangle = -E_0/E_1|\xi\rangle$. The condition $\hat{\mathcal{F}}|z\rangle = z|z\rangle$, with $\hat{\mathcal{F}} = \hat{G}_1^{(0)} + \hat{F}_1^{(1)}$, is then imposed upon expanding $|z\rangle$ in terms of the coherent states $|\xi\rangle$

$$|z\rangle \;=\; \int \mathrm{d}\mu(\xi) \langle\xi|z\rangle |\xi\rangle \;,$$

on account of the completeness relation $\displaystyle\int \mathrm{d}\mu(\xi)|\xi\rangle\langle\xi| \equiv \mathbb{I}$, $\mathrm{d}\mu(\xi) = \mathrm{d}^2\xi/\pi$, with $\mathrm{d}^2\xi = \mathrm{d}(\mathrm{Re}\xi)\mathrm{d}(\mathrm{Im}\xi)$, being the measure in the complex ξ-plane. Since $\hat{F}_1^{(1)}|\xi\rangle = \displaystyle\sum_{n=0}^{\infty}\Delta_n(\xi;t)|n\rangle$, where the coefficients $\Delta_n(\xi;t)$ are readily calculated in view of Eqs. (21), (22), one has

$$\hat{\mathcal{F}}|z\rangle \;=\; \int \mathrm{d}\mu(\xi)\langle\xi|z\rangle \left[i\eta\xi|\xi\rangle + \int \mathrm{d}\mu(\xi') \sum_{n=0}^{\infty} \Delta_n(\xi;t) c_n^*(\xi') \right] |\xi'\rangle \;. \tag{38}$$

In Eq. (38), the expansion of $|n\rangle$ in terms of the coherent states $|\xi'\rangle$, $|n\rangle = \displaystyle\int \mathrm{d}\mu(\xi')\langle\xi'|n\rangle|\xi'\rangle$, with $\langle\xi'|n\rangle = c_n^*(\xi')$, has been utilized. Left-multiplying $\hat{\mathcal{F}}|z\rangle = z|z\rangle$ by $\displaystyle\int \mathrm{d}\mu(\xi'')\langle\xi''|$ and resorting to Eq. (38), one finds

$$z \;=\; i\eta\xi + \int \mathrm{d}\mu(\xi') \mathcal{K}(\xi,\xi';t) \;, \tag{39}$$

where $\mathcal{K}(\xi,\xi';t)$ denotes the kernel

$$\mathcal{K}(\xi,\xi';t) \;=\; \sum_{n=0}^{\infty} \Delta_n(\xi;t) c_n^*(\xi') \;.$$

The time-dependent shift (39) of the eigenvalue $-E_0/E_1 = i\eta\xi$ of the operator $\hat{G}_1^{(0)}$ implies that also the values of the Lamb-Dicke parameter, possibly giving rise to a singularity, are shifted from the pathological value to a close one, most likely not singular.

VII. CONCLUSIONS

To summarize, we have analyzed the properties of the NLCS, which correspond to the experimental scheme of laser-driven trapped ions. Upon verifying that they can be defined as both Barut-Girardello states (eigenstates of annihilation operator) and minimum-uncertainty states, we have carried out a mathematical investigation whose main conclusions, in view of physical applications, are:

- For single-boson NLCS, if the squared Lamb-Dicke parameter happens to be a root of a given Laguerre polynomial $L^{(0)}_{\beta-1}(\eta^2)$ both the coherence and the minimum-uncertainty properties do not remain valid; however, the latter feature is still maintained over a wide range of typical values of η if β is not exceedingly high (i.e., one does not consider very excited vibrational levels); if the squared Lamb-Dicke parameter happens to be a root of a given Laguerre polynomial $L^{(1)}_{\mu-1}(\eta^2)$, the very existence of the single-boson NLCS is compromised, due to the oscillating behaviour of the defining series;
- the general construction which leads to the k-boson NLCS is naturally affected by the same kind of pathologies;
- we have proposed to relax the RWA conditions to get around the above difficulties and have discussed, for the single-boson NLCS, first-order corrections of the known theory;
- from an experimental point of view our results suggest that, due to the finite accuracy in fixing the parameter η when preparing single-boson and k-boson NLCS for actual experiments, one should be cautiously far from the singularities analyzed in this paper in order to maintain the RWA approach.

Acknowledgements

Two of the authors (M. G. and M. R.) would like to acknowledge the financial support of Regione Piemonte through the Research Project E14.

[1] W. Vogel and D. G. Welsch, *Quantum Optics* (Wiley-VCH, Weinheim, 2006).
[2] V. V. Dodonov, M. A. Marchiolli, Ya. A. Korennoy, V. I. Man'ko, and Y. A. Moukhin, Phys. Rev. A **58**, 4087 (1998).
[3] W. Wu and L. Wu, J. Opt. B **5**, 364 (2003).
[4] I. Jex and V. Bužek, J. Mod. Opt **40**, 771 (1993).
[5] K. Fujii, e-print arXiv: quant-ph/0112090.
[6] S. Sivakumar, J. Opt. B: Quantum Semiclass. Opt. **2**, R61 (2000).
[7] R. L. de Matos Filho and W. Vogel, Phys. Rev. A **54**, 4560 (1996).
[8] F. A. Raffa and M. Rasetti, Laser Phys. **16**, 1486 (2006).
[9] F. A. Raffa and M. Rasetti, Int. J. Quant. Inf. **5**, 229 (2007).
[10] A. Napoli and A. Messina, Eur. Phys. J. D **5**, 441 (1999).
[11] J. Katriel and A. Solomon, Phys. Rev. A **49**, 5149 (1994).
[12] V. I. Man'ko, G. Marmo, E. C. G. Sudarshan, and F. Zaccaria, Phys. Scripta **55**, 528 (1997).
[13] M. K. Tavassoly and A. Parsaiean, J. Phys. A: Math. Theor. **40**, 9905 (2007).
[14] B. R. Lavoie and H. de Guise, J. Phys. A: Math. Theor. **40**, 2825 (2007).
[15] C. C. Gerry, Opt. Comm. **91**, 247 (1992).
[16] M. Genovese and C. Novero, Phys. Rev. A **61**, 032102 (2000).
[17] J. S. Neergard-Nielsen, B. Melholt Nielsen, C. Hettich, K. Mølmer, and E. S. Polzik, Phys. Rev. Lett. **97**, 083604 (2006).
[18] A. Ourjumtsev, R. Tualle-Brouri, J. Laurat, and P. Grangier, Science **312**, 83 (2006).
[19] N. Ba An, Chinese J. Phys. **39**, 594 (2001), e-print arXiv: quant-ph/0103077.
[20] H. Moya-Cessa, A. Vidiella-Barranco, J. A. Roversi, D. S. Freitas, and S. M. Dutra, Phys. Rev. A **59**, 2518 (1999).
[21] K. Mølmer and A. Sørensen, Phys. Rev. Lett. **82**, 1835 (1999).
[22] D. Wang, T. Hansson, Å. Larson, H. O. Karlsson, and J. Larson, Phys. Rev. A **77**, 053808 (2008).
[23] V. Man'ko, G. Marmo, A. Porzio, S. Solimeno, and F. Zaccaria, Phys. Rev. A **62**, 053407 (2000).
[24] Z. Kis, W. Vogel, and L. Davidovich, Phys. Rev. A **64**, 033401 (2001).
[25] R. L. de Matos Filho and W. Vogel, Phys. Rev. Lett. **76**, 608 (1996).
[26] C. A. Blockey and D. F. Walls, Phys. Rev. A **47**, 2115 (1993).
[27] J. Črnugelj, M. Martinis, and V. Mikuta-Martinis, Phys. Rev. A **50**, 1785 (1994).
[28] L. F. Wei, S. Y. Liu, and X. L. Lei, Phys. Rev. A **65**, 062316 (2002).
[29] C. C. Gerry, S.-C. Gou, and J. Steinbach, Phys. Rev. A **55**, 630 (1997).
[30] A. O. Barut and L. Girardello, Commun. Math. Phys. **21**, 41 (1971).

[31] For example $L_{\hat{n}}^{(\alpha)}(x) = \dfrac{1}{2\pi i} \oint_C \dfrac{e^{-xz/(1-z)}\, z^{-\hat{n}}}{(1-z)^{\alpha+1}} \dfrac{\mathrm{d}z}{z}$, where C is a closed contour centered at the origin of the complex plane, whose radius $|z|$ is < 1.
[32] M. Abramowitz and I. A. Stegun (Eds.), *Handbook of Mathematical Functions* (Dover, New York, 1970).
[33] P. Zanardi and M. Rasetti, Phys. Rev. Lett. **79**, 3306 (1997).
[34] A. Perelomov, *Generalized Coherent States and their Applications* (Springer, Berlin, 1986).
[35] G. Szegö, *Orthogonal Polynomials* (4th ed., American Mathematical Society, Providence, Rhode Island, 1975).